

Prenatal Exposure to Phthalates and Newborn Telomere Length: A Birth Cohort Study in Wuhan, China

Lulu Song,^{1,2} Bingqing Liu,^{1,2} Mingyang Wu,^{1,2} Lina Zhang,^{1,2} Lulin Wang,^{1,2} Bin Zhang,³ Chao Xiong,³ Yuanyuan Li,² Zhongqiang Cao,³ Youjie Wang,^{1,2} and Shunqing Xu²

¹Department of Maternal and Child Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

²Key Laboratory of Environment and Health, Ministry of Education & Ministry of Environmental Protection, and State Key Laboratory of Environmental Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

³Wuhan Children's Hospital (Wuhan Maternal and Child Healthcare Hospital), Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

BACKGROUND: Telomere length (TL) is a marker of biological aging and is inversely related to aging-related diseases. The setting of TL at birth may have important implications for lifelong telomere dynamics; however, its determinants remain poorly understood.

OBJECTIVES: The purpose of our study was to explore the relationships between prenatal exposure to phthalates and umbilical cord blood TL.

METHODS: A total of 762 mother–newborn pairs were recruited from a birth cohort study performed between November 2013 and March 2015 in Wuhan, China. Relative cord blood TL was measured using quantitative real-time polymerase chain reaction. Six phthalate metabolites were measured in urine samples acquired from pregnant women during the three trimesters. Multiple informant models were applied to estimate the associations between prenatal exposure to phthalates and cord blood TL and to evaluate potential windows of vulnerability.

RESULTS: Exposure to mono-ethyl phthalate (MEP), mono(2-ethyl-5-carboxypentyl) phthalate (MECPP), mono(2-ethyl-5-oxohexyl) phthalate (MEOHP), mono(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP), mono-butyl phthalate (MBP), and di(2-ethylhexyl) phthalate (Σ DEHP) during the first trimester were inversely related to cord blood TL. In addition, we observed a female-specific association between maternal exposure to MEP during the first trimester and cord blood TL ($p_{\text{sex-int}} = 0.03$). The associations between maternal exposure to MECPP, MEHHP, MEOHP, and Σ DEHP during the first trimester and cord blood TL were consistent between males and females (all $p_{\text{sex-int}} > 0.10$).

CONCLUSION: This prospective study demonstrated that prenatal exposure to some phthalate metabolites were associated with shorter cord blood TL. Our results, if confirmed in other populations, may provide more evidence of adverse health outcomes of phthalate exposure and support the hypothesis that the intrauterine environment may be one of the major determinants for newborn TL. <https://doi.org/10.1289/EHP4492>

Introduction

Phthalates are widely used in plasticizers or solvents in consumer products such as cosmetics, toys, plastics, food packing, and building materials (Koch and Calafat 2009; Schettler 2006). Because phthalates are noncovalently bound to the products, they are easily released into the external environment and exposure may occur in humans through ingestion, inhalation, and dermal absorption (Schettler 2006; Wittassek et al. 2011). Food intake is the main route of high-molecular-weight phthalates (high-MWPs), whereas personal care products are the major source of low-molecular-weight phthalates (low-MWPs) (Rudel et al. 2011; Wormuth et al. 2006). Once absorbed, phthalates are quickly metabolized and excreted in the urine, where they can be measured. Phthalate metabolites have been consistently detected in urine from the general population and from pregnant women worldwide (Benjamin et al. 2017; Johns et al. 2015).

Due to the ubiquitous presence of phthalate and potential health consequences after phthalate exposure, there has been increasing concern, especially in susceptible populations such as pregnant

women and fetuses. Phthalate metabolites can cross the placenta in humans and have been detected in cord blood (Latini et al. 2003; Mose et al. 2007). A growing body of literature has reported that prenatal phthalate exposures were related to adverse perinatal outcomes such as preeclampsia, preterm birth, pregnancy loss, and fetal retarded growth (Cantonwine et al. 2016; Casas et al. 2016; Ferguson et al. 2014c; Mu et al. 2015). Although phthalates have known endocrine-disrupting properties (Diamanti-Kandarakis et al. 2009; Halden 2010), there is also evidence from human studies that phthalate metabolites are related to increased biomarkers of oxidative stress (e.g., 8-isoprostane and 8-hydroxydeoxyguanosine) and inflammation [e.g., tumor necrosis factor alpha (TNF- α), interleukin-6 (IL-6), IL-10, and IL-1 β] in pregnant women (Ferguson et al. 2012, 2014a, 2015; Holland et al. 2016). Oxidative stress and chronic inflammation are considered potential pathways by which exposure to phthalates may influence adverse health effects (Ferguson et al. 2017; Holland et al. 2016).

Telomeres consist of noncoding repetitive sequences of DNA (TTAGGG) and nucleoproteins that together form a “cap” at the end of chromosomes and protect the integrity of the genomic content (Blackburn 1991). Telomere length (TL) shortens with each cell division and reflects biological aging (Benetos et al. 2001). Both high oxidative stress and inflammation have been linked to TL shortening. Telomeres with guanine-rich sequences are more sensitive to oxidative insult (Houben et al. 2008; Kawanishi and Oikawa 2004). Elevated inflammatory activity could accelerate leukocyte telomere shortening by promoting cell turnover (Jurk et al. 2014) as well as by inducing reactive oxygen species that damage telomeric DNA via oxidative stress (Jaiswal et al. 2000). Shorter TL has been related to aging-related diseases such as cardiovascular disease, type 2 diabetes, cancer, and all-cause mortality (Haycock et al. 2014; Mons et al. 2017; Willeit et al. 2010; Zhao et al. 2013).

Although studies on TL have been mainly conducted in adults, TL in early life is being increasingly considered as critical

Address correspondence to Youjie Wang, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Hangkong Rd., Wuhan 430030, Hubei, China. Telephone: (86)27-83691198. Email: Wangyoujie@mails.tjmu.edu.cn

Supplemental Material is available online (<https://doi.org/10.1289/EHP4492>).

The authors declare they have no actual or potential competing financial interests.

Received 19 September 2018; Revised 1 August 2019; Accepted 5 August 2019; Published 26 August 2019.

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

to an individual's long-term health and disease risk (Fairlie et al. 2016; Heidinger et al. 2012; Moreno-Palomo et al. 2014). TL at birth represents an individual's initial setting of TL, which means that TL in adulthood is determined by newborn TL and subsequent attrition (Benetos et al. 2013; Hjelmborg et al. 2015; Sabharwal et al. 2018). Thus, factors that influence initial or newborn TL may be the potential determinants of lifetime health. Several studies have been performed to explore the determinants of newborn TL, such as maternal prepregnancy body mass index (BMI) (Martens et al. 2016), maternal psychosocial stress (Marchetto et al. 2016), maternal smoking (Mirzakhani et al. 2017), maternal exposure to polycyclic aromatic hydrocarbons (PAHs) (Perera et al. 2018), maternal particulate matter exposure (Martens et al. 2017), maternal residential traffic exposure (Bijns et al. 2015), maternal heavy metal exposure (Wai et al. 2018), and maternal exposure to perfluoroalkyl and polyfluoroalkyl substances (Liu et al. 2018). However, studies regarding the effect of prenatal phthalate exposure on newborn TL are lacking. To date, only one cross-sectional study reported positive associations between urinary phthalates and TL in adults (Scinicariello et al. 2016).

Given that *in utero* life is considered to be a critical period of vulnerability in the early programming of subsequent health (Barker 1990), uncovering the associations of prenatal exposure to phthalate metabolites with newborn TL may provide new insights into the pathways of phthalate-related disorders. Therefore, we explored the associations of prenatal exposure to phthalate metabolites with newborn TL in a birth cohort study.

Methods

Study Population

The birth cohort study was conducted at the Wuhan Children's Hospital (Wuhan Maternal and Child Healthcare Hospital) between November 2013 and March 2015. Briefly, 762 pregnant women were enrolled in the study during their first prenatal visit in Wuhan city, Hubei province, China. We recruited the participants using the following inclusion criteria: *a*) resident of Wuhan city, *b*) <16 weeks of pregnancy with a singleton gestation at recruitment, and *c*) planning to take prenatal care and delivery in the study hospital. In the present study, we excluded 16 participants due to unavailability of cord blood or ineligible DNA quality. The final analysis included 746 mother–newborn pairs. Of the eligible pregnant women, the number of women with urine samples was 746, 746, and 743 for the first, second, and third trimesters, respectively.

The study protocol was approved by the ethics committees of Tongji Medical College, Huazhong University of Science and Technology, and the Wuhan Children's Hospital (Wuhan Maternal and Child Healthcare Hospital). All women provided written informed consent at enrollment.

Cord Blood Collection and TL Measurements

Umbilical cord blood was drawn immediately at delivery. Blood samples were centrifuged and stored at -80°C until DNA extraction. DNA was extracted from the leukocytes of umbilical cord blood using the Wizard[®] Genomic DNA Purification (Promega Corporation). To ensure good quality DNA samples, DNA concentration and purity were determined using a NanoDrop Spectrophotometer (ND-1000; Thermo Fisher Scientific), and those with the ratio of A260/280 ranging from 1.8 to 2.0 were considered eligible. Relative cord blood TL was determined based on the ratio of the telomere (T) repeat copy number to the single-copy gene (S) copy number (T/S ratio) for each indi-

vidual using the quantitative real-time polymerase chain reaction (qPCR) method, as described previously (Cawthon 2009; Qu et al. 2015). For the telomeres, the forward and reverse primers were 5'-ACACTAAGGTTTGGGTTTGGGTTTGGGTTTGGGTTAGTGT-3' and 5'-TGTTAGGTATCCCTATCCCTATCCCTATCCCTATCCCTAACA-3' (Cawthon 2009). For the reference gene [human beta-globin (hbg)], the forward and reverse primers were 5'-GTGCACCTGACTCCTGAGGAGA-3' and 5'-CCTTGATACCAACCTGCCCAG-3' (Qu et al. 2015). We randomly selected 50 DNA samples from our study population and pooled them as the reference DNA, which was used to obtain a standard curve, ranging from 104 to 0.4 ng/ μL ($R^2 > 0.99$). Each PCR was conducted using a 1.0- μL DNA sample (10 ng/ μL) in a final volume of 10 μL per reaction. The telomere reaction mixture contained KAPA SYBR[®] FAST qPCR Kit Master MIX 1 \times (KAPA Biosystems; 5 μL /reaction), forward (0.27 μL /reaction–270 nM) and reverse (0.90 μL /reaction–900 nM) primers, and RNase-free water (2.83 μL /reaction). The single-copy gene reaction mixture contained KAPA SYBR[®] FAST qPCR Kit Master MIX 1 \times (KAPA Biosystems; 5 μL /reaction), forward (0.2 μL /reaction–200 nM) and reverse (0.2 μL /reaction–200 nM) primer, and RNase-free water (3.6 μL /reaction). The PCR thermal cycling conditions were set up as follows: 2 min at 50°C and 3 min at 95°C for activation of the DNA polymerase, followed by 40 cycles consisted of 3 s at 95°C for denaturation and 30 s at 60°C for annealing/extension. All samples were run in triplicate in 384-well plates using the ViiA[™] 7 Dx Real-Time PCR System (Applied Biosystems). The intra-run and inter-run coefficients of variation for the TL measurements were 3.0% and 4.1%, respectively.

Urine Sample Collection and Phthalate Metabolite Measurements

We collected spot urine samples of the pregnant women at 13.0 ± 1.1 , 23.6 ± 3.2 , and 36.0 ± 3.3 weeks of gestation. The gestational ages of urine collection roughly reflect the first, second, and third trimesters. All urine samples were collected in polypropylene tubes and frozen at -20°C for further analysis.

We analyzed six phthalate metabolites in urine samples, including mono-ethyl phthalate (MEP), mono(2-ethyl-5-oxohexyl) phthalate (MEOHP), mono(2-ethylhexyl) phthalate (MEHP), mono(2-ethyl-5-hydroxyhexyl) phthalate (MEHHP), mono(2-ethyl-5-carboxypentyl) phthalate (MECPP), and mono-butyl phthalate (MBP). In addition to measuring the individual phthalate metabolites, the total concentration of di(2-ethylhexyl) phthalate (ΣDEHP) was estimated using the sum of the molar concentrations of MECPP, MEHP, MEHHP, and MEOHP. We analyzed phthalate metabolites in urine samples using solid-phase extraction coupled with ultra-performance liquid chromatography–tandem mass spectrometry (UPLC-MS/MS). The details of the analytical procedure have been previously described (Zhu et al. 2016, 2018). The limits of detection (LODs) were 0.01 ng/mL for MECPP, 0.05 ng/mL for MEOHP and MEHHP, and 0.1 ng/mL for MEP, MBP, and MEHP. Urinary phthalate metabolite concentrations <LOD were determined as the LOD divided by the square root of 2.

To adjust for variations in urine dilution, we corrected urinary phthalate concentrations by urinary specific gravity (SG) based on the following equation: $P_{\text{SG}} = P[(1.012 - 1)/(SG - 1)]$, where P_{SG} is the SG-corrected phthalate concentration, P is the observed phthalate concentration, the value 1.012 is the median SG in this study population, and SG is the specific gravity of the individual urine samples. SG was measured using a refractometer (Atago PAL-10S; Atago) while preparing urine samples for phthalate analysis. The SG-corrected phthalate concentrations were used for statistical analysis.

Covariates

Information on sociodemographic and lifestyle characteristics was obtained through standard questionnaires administered by trained nurses at the first prenatal care visit and included maternal age, educational level, and passive smoking during pregnancy. Passive smoking during pregnancy was defined as exposure to secondhand smoke from family members in the household or colleagues at the workplace during pregnancy (Vardavas et al. 2016). Information on mothers' reproductive history, hypertensive disorders of pregnancy, gestational diabetes, and infants' sex and birth date was retrieved from medical records. Gestational age (weeks) was determined based on the last menstrual period. We defined gestational age <37 weeks as preterm birth. Prepregnancy weight was self-reported during the first prenatal care visit and height was measured using a stadiometer. The prepregnancy BMI was determined by dividing self-reported prepregnancy weight (in kilograms) by height squared (in meters squared).

Statistical Analysis

Continuous data were expressed as the mean \pm standard deviation (SD) for normal distributions, geometric mean (interquartile range) for skewed distributions, or n (%) for categorical data. Urinary phthalate levels and cord blood TL were natural log-transformed to improve the normality of the distributions. Correlations between individual phthalate metabolites and SG were estimated using Pearson correlation coefficients or Partial Pearson correlation coefficients that adjusted for trimester. Reproducibility of urinary phthalate concentrations across the different three trimesters was determined by computing the intraclass correlation coefficients (ICCs) according to random intercept linear mixed models. The ICC was defined as the ratio of the between-person variance to the sum of within- and between-person variances.

Multiple informant models were applied to explore the associations between repeated prenatal urinary phthalate concentrations and newborn TL (Sánchez et al. 2011). The model treated individual phthalate concentrations in different trimesters (or windows) as informants and jointly estimated the difference in newborn TL for each doubling of urinary phthalate concentrations in each trimester, using a nonstandard version of generalized estimating equations. In this model, each exposure window did not adjust for urinary phthalate concentrations in other windows, but adjusted for the same covariates. The multiple informant model also provided a way to test differences in associations between urinary phthalate concentrations and newborn TL across the three trimesters. The null hypothesis for the test was that the phthalate coefficients were equal across the three trimesters.

To capture potential nonlinear relationships, we also used multiple informant models to assess the associations between quartiles of prenatal urinary phthalate concentrations and cord blood TL, with the lowest quartile as the reference. Quartiles of urinary phthalate concentrations were categorized based on trimester-specific cut points. Tests for linear trend across quartiles were carried out by assigning the median value of each quartile as a continuous variable. To enhance the interpretation of regression models consisting of ln-transformed exposure and/or outcome variables, we presented all regression coefficients and 95% confidence intervals (CIs) as percent change and 95% CI in cord blood TL for a doubling or quartiles of urinary phthalate concentrations. Multivariate models were adjusted for maternal age (continuous), prepregnancy BMI (continuous), educational level (junior high school or below, high school, college or above),

parity (primiparous, multiparous), passive smoking during pregnancy (yes, no), and infant sex (male, female).

Because previous investigations have suggested a sex-specific relationship between prenatal exposure to phthalates and fetal development, we explored whether infant sex modified the association between prenatal exposure to phthalates and newborn TL. Stratum-specific p -values for interaction between each phthalate metabolite and infant sex were estimated separately for each trimester using the Wald test (Kaufman and MacLehose 2013), with a $p < 0.1$ considered significant. We also conducted a series of sensitivity analyses with additional adjustment for gestational age at urine collection, excluding mothers with hypertensive disorders of pregnancy and gestational diabetes, and excluding preterm infants. Moreover, we fitted separate linear regression models to explore the relationships between prenatal exposure to phthalates and cord blood TL.

All statistical analyses were carried out using SAS (version 9.4; SAS Institute Inc.). A two-sided $p < 0.05$ was considered statistically significant.

Results

The characteristics of the 746 mother–newborn pairs are presented in Table 1. The mean prepregnancy BMI and maternal age of mothers were 20.8 ± 2.8 kg/m² and 28.6 ± 3.3 y, respectively. Among the 746 mothers, 589 (79.0%) were well educated, 244 (32.7%) were passively exposed to cigarette smoking, and 644 (86.3%) were primiparous. Approximately 19 (2.5%) and 53 (7.1%) of mothers were diagnosed with hypertensive disorders of pregnancy and gestational diabetes, respectively. Of the newborns, 383 (51.3%) were males. The rate of preterm birth was 2.4%. The characteristics of mothers carrying male infants were similar to those carrying female infants.

The detection rate, distributions, and variability (i.e., the ICCs) of SG-corrected urinary phthalate metabolites in each trimester are shown in Table 2. The detection rate of phthalate metabolites ranged from 87.0% to 99.9%. MBP had the highest median concentration in each trimester, followed by MEP, MECPP, MEHHP, MEOHP, and MEHP. The ICCs for urinary phthalate concentrations during the three trimesters ranged from 0.19 to 0.36, with the lowest ICC for MEHHP and the highest for MEP.

The detection rate and distributions of SG-corrected urinary phthalate metabolites during each trimester stratified by infant sex are shown in Table S1. The Pearson correlation coefficients between SG-corrected urinary phthalate metabolites ranged from 0.08 to 0.91 for the first trimester, 0.08 to 0.92 for the second trimester, and 0.05 to 0.96 for the third trimester, with the highest Pearson correlation coefficient between MEHHP and MEOHP in each trimester (see Tables S2–S4). The partial Pearson correlation coefficients of SG-corrected urinary phthalate metabolites adjusted for trimester ranged from 0.07 to 0.93 (see Table S5).

Table 3 displays the relationships of prenatal exposure to phthalate metabolites with cord blood TL. In the multivariate-adjusted models, each doubling of maternal urinary MECPP, MEHHP, and MEOHP during the first trimester were associated with 2.38% (95% CI: -4.22% , -0.51%), 2.53% (95% CI: -4.38% , -0.64%), and 2.90% (95% CI: -4.76% , -1.01%) shorter cord blood TL, respectively, with consistent inverse associations for the second and third trimesters. In addition, each doubling of maternal urinary Σ DEHP during the first trimester was associated with a 3.25% (95% CI: -5.26% , -1.20%) decrease in cord blood TL. The pattern was similar for the second trimester, whereas a positive association was observed for the third trimester.

Table 1. Characteristics of mother–newborn pairs, Wuhan, China, 2013–2015 (*n* = 746).

Characteristic	Total (<i>n</i> = 746)	Mothers carrying male infants (<i>n</i> = 383)	Mothers carrying female infants (<i>n</i> = 363)	<i>p</i> -Value
Maternal age (y)	28.6 ± 3.3	28.7 ± 3.5	28.5 ± 3.2	0.24
Prepregnancy BMI (kg/m ²)	20.8 ± 2.8	21.0 ± 2.9	20.8 ± 2.7	0.27
Educational level [<i>n</i> (%)]				0.77
Junior high school or below	42 (5.6)	23 (6.0)	19 (5.2)	
High school	115 (15.4)	56 (14.6)	59 (16.3)	
College or above	589 (79.0)	304 (79.4)	285 (78.5)	
Passive smoking [<i>n</i> (%)]				0.56
Yes	244 (32.7)	129 (33.7)	115 (31.7)	
No	502 (67.3)	254 (66.3)	248 (68.3)	
Gestational diabetes [<i>n</i> (%)]				0.82
Yes	53 (7.1)	28 (7.3)	25 (6.9)	
No	693 (92.9)	355 (92.7)	338 (93.1)	
Hypertensive disorders of pregnancy [<i>n</i> (%)]				0.30
Yes	19 (2.5)	12 (3.1)	7 (1.9)	
No	727 (97.5)	371 (96.9)	356 (98.1)	
Parity [<i>n</i> (%)]				0.10
Primiparous	644 (86.3)	323 (84.3)	321 (88.4)	
Multiparous	102 (13.7)	60 (15.7)	42 (11.6)	
Preterm birth (gestational age <37 weeks) [<i>n</i> (%)]				0.91
Yes	18 (2.4)	9 (2.3)	9 (2.5)	
No	728 (97.6)	374 (97.7)	354 (97.5)	
Cord blood telomere length (T/S ratio)	0.75 (0.56–0.95)	0.75 (0.55–0.97)	0.74 (0.57–0.93)	0.98

Note: Continuous variables are presented as mean ± SD (normally distributed) or geometric mean with 25–75th percentile (not normally distributed); categorical variables are expressed as *n* (%). Data were complete for all variables.

We also explored the associations between quartiles of prenatal exposure to phthalate metabolites and cord blood TL (see Table S6). Compared with maternal urinary MEP, MECPP, MEHHP, and MEOHP during the first trimester in the lowest quartile, those in the highest quartile were associated with 11.12% (95% CI: −17.95%, −3.73%), 11.64% (95% CI: −18.42%, −4.30%), 9.30% (95% CI: −16.26%, −1.74%), and 11.78% (95% CI: −18.56%, −4.43%) shorter cord blood TL, with similar patterns of associations for the second and third trimesters. Moreover, mothers in the highest quartile of urinary MBP and ΣDEHP during the first trimester had 8.43% (95% CI: −15.46%, −0.83%) and 11.98% (95% CI: −18.75%, −4.64%) shorter cord blood TL than in the lowest quartile, respectively. The inverse association of ΣDEHP was consistent for the second trimester, whereas positive associations of MBP and ΣDEHP were observed for the third trimester. Maternal urinary MEHP was not significantly associated with cord blood TL, regardless of trimester (Table 3; see also Table S6).

Analysis stratified by infant sex is presented in Table 4. Maternal urinary MEP during the first trimester was inversely associated with cord blood TL in females, whereas the association was close to the null for males (percent change, −3.07%; 95% CI: −5.20%, −0.89% and percent change, 0.24%; 95% CI:

−1.70%, 2.22%, respectively, $p_{\text{sex-int}} = 0.03$). The pattern of MEP was similar for the second trimester, whereas the inverse association was stronger in males than females for the third trimester; however, differences between males and females were not significant (all $p_{\text{sex-int}} > 0.10$). The inverse associations between maternal urinary MECPP, MEOHP, MEHHP, and ΣDEHP during the first trimester and cord blood TL were consistent between males and females (all $p_{\text{sex-int}} > 0.10$).

In the sensitivity analyses, the inverse relationships between maternal exposure to phthalate metabolites during the first trimester and cord blood TL were not materially changed with additional adjustment for gestational age at urine collection, or excluding mothers with hypertensive disorders of pregnancy and gestational diabetes (*n* = 68), or excluding preterm infants (*n* = 18) (see Table S7). In separate linear regression models, the relationships between prenatal phthalate exposure and cord blood TL were similar to those in the main analyses (see Table S8).

Discussion

In this study, we observed that exposure to MEP, MECPP, MEOHP, MEHHP, MBP, and ΣDEHP during the first trimester were inversely related to cord blood TL. The stratified analysis

Table 2. Distributions and intraclass correlation coefficients of maternal urinary phthalate metabolites according to trimester, Wuhan, China, 2013–2015.

Phthalate metabolites (ng/mL)	Percentile																ICC
	>LOD (%)			GM			25th			50th			75th				
	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd		
MEP	97.8	98.9	98.9	10.12	9.10	10.18	5.04	4.13	4.72	9.72	8.07	9.80	20.28	20.41	22.54	0.36	
MECPP	99.5	99.9	99.7	9.52	8.53	9.86	5.68	5.00	6.29	9.24	8.30	9.83	15.95	13.48	15.56	0.24	
MEHHP	99.6	99.9	99.9	6.53	5.14	6.12	3.64	2.66	3.62	6.18	4.94	6.14	11.43	8.67	10.01	0.19	
MEOHP	99.6	99.9	99.9	4.81	4.13	5.03	2.68	2.17	3.06	4.80	4.25	4.99	8.23	6.87	8.09	0.24	
MBP	98.1	98.3	98.5	64.86	54.56	86.91	34.17	25.76	41.11	65.44	58.50	95.58	142.7	133.2	207.4	0.22	
MEHP	89.2	88.0	87.0	2.35	1.88	1.85	1.08	0.89	0.81	2.81	2.18	2.24	5.68	4.85	4.98	0.32	
ΣDEHP (nmol/L)	—	—	—	86.33	71.57	85.24	49.37	39.39	50.97	80.52	68.50	81.82	140.0	113.8	130.3	0.21	

Note: The ICC was derived by dividing the between-person variance by the sum of the between- and within-person variances. Values range from 0 (no reproducibility of the same measurement within a subject) to 1 (perfect reproducibility). Phthalate metabolite concentrations are shown after substituting the >LOD values by LOD divided by the square root of 2. —, not applicable; DEHP, di(2-ethylhexyl) phthalate; GM, geometric mean; ICC, intraclass correlation coefficient; LOD, limit of detection; MBP, mono-butyl phthalate; MECPP, mono(2-ethyl-5-carboxypentyl) phthalate; MEHHP, mono(2-ethyl-5-hydroxyhexyl) phthalate; MEHP, mono(2-ethylhexyl) phthalate; MEOHP, mono(2-ethyl-5-oxohexyl) phthalate; MEP, mono-ethyl phthalate; ΣDEHP, molecular sum of MECPP, MEHP, MEHHP, and MEOHP (nmol/L).

Table 3. Associations between prenatal exposure to phthalate metabolites and cord blood telomere length, Wuhan, China, 2013–2015.

Phthalate metabolites	1st trimester (n = 746)		2nd trimester (n = 746)		3rd trimester (n = 743)		<i>p</i> _{int} ^a
	Percent change (95% CI)	<i>p</i> -Value	Percent change (95% CI)	<i>p</i> -Value	Percent change (95% CI)	<i>p</i> -Value	
MEP	−1.09 (−2.53, 0.37)	0.14	−0.48 (−2.01, 1.09)	0.55	−0.92 (−2.41, 0.60)	0.23	0.74
MECPP	−2.38 (−4.22, −0.51)	0.01	−0.72 (−2.73, 1.33)	0.49	−0.87 (−2.96, 1.27)	0.42	0.31
MEHHP	−2.53 (−4.38, −0.64)	0.009	−1.25 (−3.18, 0.72)	0.21	−0.29 (−2.50, 1.97)	0.80	0.29
MEOHP	−2.90 (−4.76, −1.01)	0.003	−1.12 (−3.04, 0.83)	0.26	−0.51 (−2.65, 1.67)	0.64	0.16
MBP	−1.13 (−2.52, 0.28)	0.12	0.57 (−0.79, 1.95)	0.41	1.19 (−0.22, 2.62)	0.10	0.04
MEHP	−1.18 (−2.44, 0.11)	0.07	−0.30 (−1.60, 1.02)	0.65	0.88 (−0.35, 2.14)	0.16	0.02
ΣDEHP	−3.25 (−5.26, −1.20)	0.002	−0.72 (−2.73, 1.34)	0.49	0.63 (−1.70, 3.01)	0.60	0.02

Note: All models were adjusted for maternal age, prepregnancy BMI, parity, educational level, passive smoking during pregnancy, and infant sex. BMI, body mass index; CI, confidence interval; DEHP, di(2-ethylhexyl) phthalate; MBP, mono-butyl phthalate; MECPP, mono(2-ethyl-5-carboxypentyl) phthalate; MEHHP, mono(2-ethyl-5-hydroxyhexyl) phthalate; MEHP, mono(2-ethylhexyl) phthalate; MEOHP, mono(2-ethyl-5-oxohexyl) phthalate; MEP, mono-ethyl phthalate; *p*_{int}, *p*-value for interaction; ΣDEHP, molecular sum of MECPP, MEHP, MEHHP, and MEOHP (nmol/L).

^aScore test of homogeneity of regression coefficients.

suggested that maternal urinary MEP during the first trimester was associated with shorter cord blood TL in female infants, whereas the association was close to the null for male infants. The inverse associations between maternal urinary MECPP, MEOHP, MEHHP, and ΣDEHP during the first trimester and cord blood TL were consistent between male and female infants.

Phthalate metabolites were detectable in maternal urine samples in our study, owing to the ubiquitous presence of phthalates. The concentrations of some phthalate metabolites (e.g., MEHHP, MEHP, and MEOHP) in our study were comparable with those in other studies of Chinese pregnant women (Gao et al. 2017; Zhang et al. 2018; Zhu et al. 2018). However, most concentrations of urinary phthalate metabolites (MEHHP, MECPP, MEHP, and MEOHP) among pregnant women from the United States (Soomro et al. 2018), Mexico (Téllez-Rojo et al. 2013), Spain (Casas et al. 2016), France (Philippat et al. 2012), and Greece

(Myridakis et al. 2015) were higher than those reported in the present study. Compared with pregnant women from Canada (Arbuckle et al. 2014) and Denmark (Tefre de Renzy-Martin et al. 2014), similar concentrations of urinary MECPP, MEHP, MEOHP, and MEHHP and lower concentrations of MEP were shown in our study. The differences in phthalate exposure levels in different regions may result from variations in lifestyle behaviors, use of consumer products, and living environments. In our study, we observed that ICCs for urinary phthalate concentrations ranged from 0.19 to 0.36, which indicated a high variability across the whole pregnancy. ICCs for MEOHP and MEHP throughout pregnancy in our study were higher than those reported among studies from the United States (Ferguson et al. 2014b) and Mexico (Watkins et al. 2017a). Compared with previous studies (Ferguson et al. 2014b; Watkins et al. 2017b; Zhang et al. 2018), lower ICCs for most urinary phthalate metabolites (MEP, MECPP, MEHHP, and MBP) were shown in

Table 4. Associations between prenatal exposure to phthalate metabolites and cord blood telomere length, stratified by infant sex, Wuhan, China, 2013–2015.

Phthalate metabolites	1st trimester (n = 746)		2nd trimester (n = 746)		3rd trimester (n = 743)	
	Percent change (95% CI)	<i>p</i> -Value	Percent change (95% CI)	<i>p</i> -Value	Percent change (95% CI)	<i>p</i> -Value
MEP						
Male	0.24 (−1.70, 2.22)	0.81	−0.19 (−2.48, 2.16)	0.88	−1.37 (−3.42, 0.72)	0.20
Female	−3.07 (−5.20, −0.89)	0.006	−0.87 (−2.90, 1.21)	0.41	−0.24 (−2.41, 1.97)	0.83
<i>p</i> _{sex-int}	0.03		0.67		0.46	
MECPP						
Male	−2.05 (−4.41, 0.37)	0.10	−0.69 (−3.44, 2.13)	0.63	−1.09 (−3.91, 1.82)	0.46
Female	−3.11 (−6.08, −0.05)	0.05	−0.88 (−3.81, 2.14)	0.56	−0.27 (−3.41, 2.96)	0.87
<i>p</i> _{sex-int}	0.59		0.93		0.71	
MEHHP						
Male	−2.10 (−4.62, 0.49)	0.11	−0.74 (−3.45, 2.05)	0.60	−0.31 (−3.25, 2.73)	0.84
Female	−3.17 (−5.91, −0.36)	0.03	−1.81 (−4.54, 1.00)	0.21	0.02 (−3.35, 3.50)	0.99
<i>p</i> _{sex-int}	0.58		0.59		0.89	
MEOHP						
Male	−2.67 (−5.23, −0.04)	0.05	−0.89 (−3.53, 1.82)	0.52	−0.69 (−3.47, 2.18)	0.63
Female	−3.23 (−5.93, −0.44)	0.02	−1.40 (−4.18, 1.47)	0.34	0.11 (−3.28, 3.62)	0.95
<i>p</i> _{sex-int}	0.77		0.80		0.72	
MBP						
Male	−0.55 (−2.40, 1.34)	0.57	1.21 (−0.84, 3.30)	0.25	0.79 (−1.29, 2.90)	0.46
Female	−1.86 (−3.99, 0.31)	0.09	0.06 (−1.72, 1.87)	0.95	1.61 (−0.29, 3.56)	0.10
<i>p</i> _{sex-int}	0.37		0.41		0.57	
MEHP						
Male	−1.40 (−3.16, 0.39)	0.12	−0.88 (−2.68, 0.95)	0.34	1.12 (−0.70, 2.95)	0.23
Female	−0.84 (−2.67, 1.03)	0.38	0.48 (−1.39, 2.38)	0.62	0.67 (−1.01, 2.37)	0.44
<i>p</i> _{sex-int}	0.67		0.31		0.73	
ΣDEHP						
Male	−3.35 (−6.15, −0.48)	0.02	−0.86 (−3.64, 2.00)	0.55	0.93 (−2.18, 4.14)	0.56
Female	−3.17 (−6.05, −0.19)	0.04	−0.58 (−3.50, 2.43)	0.70	0.31 (−3.21, 3.94)	0.87
<i>p</i> _{sex-int}	0.93		0.89		0.80	

Note: All models were adjusted for maternal age, prepregnancy BMI, parity, educational level, and passive smoking during pregnancy. BMI, body mass index; CI, confidence interval; DEHP, di(2-ethylhexyl) phthalate; MBP, mono-butyl phthalate; MECPP, mono(2-ethyl-5-carboxypentyl) phthalate; MEHHP, mono(2-ethyl-5-hydroxyhexyl) phthalate; MEHP, mono(2-ethylhexyl) phthalate; MEOHP, mono(2-ethyl-5-oxohexyl) phthalate; MEP, mono-ethyl phthalate; *p*_{sex-int}, *p*-value for interaction between urinary phthalates and infant sex; ΣDEHP, molecular sum of MECPP, MEHP, MEHHP, and MEOHP (nmol/L).

the present study. The different ICCs were possibly due to differences in the consumption of personal care products, living environments, and dietary habits in different regions.

A growing body of literature has shown associations of exposure to environmental hazards with TL in adults, including metals [cadmium (Zota et al. 2015), lead (Wu et al. 2012), and arsenic (Li et al. 2012)], air pollution [black carbon (McCracken et al. 2010) and particulate matter (Hou et al. 2012; Pieters et al. 2016)], persistent organic pollutants (POPs) (Mitro et al. 2016), polychlorinated biphenyls (PCBs) (Callahan et al. 2017), and benzene (Bassig et al. 2014). However, evidence on the variations of newborn TL in relation to early life environmental exposures is limited. A birth cohort study of 641 mother–newborn pairs from Belgium suggested that prenatal particulate matter ≤ 2.5 μm in aerodynamic diameter ($\text{PM}_{2.5}$) exposure was associated with shorter newborn TL (Martens et al. 2017). Martens et al. (2016) also reported that prepregnancy BMI was associated with shorter newborn TL. Our study revealed a similar, but not statistically significant, association. The possible reason might be that the participants in the study by Martens et al. had a much higher average prepregnancy BMI than found in our study. In addition, Perera et al. (2018) found that prenatal exposure to PAHs from coal burning was inversely related to cord blood TL. Overall, our findings and those from previous studies suggest that newborn TL may be influenced by multiple environmental stressors with small individual effects. To date, the relationships between prenatal exposure to phthalates and newborn TL have not been reported. Only a cross-sectional study based on data from the National Health and Nutrition Examination Survey (NHANES) 1999–2002 ($n = 2,742$) had evaluated the relationships between urinary phthalate metabolites [MEP, MBP, MEHP, and mono-benzyl phthalate (MBzP)] and leukocyte TL in adults ≥ 20 y of age (Scinicariello et al. 2016). Contrary to our findings, Scinicariello et al. (2016) found that urinary MEHP was related to longer TL among all subjects and that urinary MBP and MBzP were related to longer TL only among people ≥ 60 y of age. Given that the telomere dynamics in early life were different from those in later life (Frenck et al. 1998; Turner et al. 2019), the difference in research subjects (newborn vs. adult) was expected to be the major contributor to the discrepancies between the two studies. In addition, differences in study design (longitudinal vs. cross-sectional) as well as exposure time period, exposure concentration, and exposure duration might be reasons for the inconsistent findings.

Although the potential mechanisms underlying the relationships between maternal exposure to phthalates and cord blood TL are yet to be elucidated, oxidative stress and chronic inflammation are two plausible hypotheses. Exposure to phthalates has been associated with increased oxidative stress among pregnant women (Ferguson et al. 2014a, 2015). Because of the high guanine content of the telomere sequence, telomeres are more vulnerable to damage induced by oxidative stress (Houben et al. 2008; Kawanishi and Oikawa 2004). Moreover, telomeric DNA is deficient in the repair of single-strand breaks, compared with genomic DNA; oxidative stress can induce single-strand breaks, resulting in increased telomere shortening (Houben et al. 2008; Kawanishi and Oikawa 2004). Exposure to phthalate metabolites during pregnancy could also affect newborn telomere biology through chronic inflammation. Evidence suggests that chronic inflammation is related to increasing amounts of leukocytes and their more rapid expenditure, which requires a higher rate of cell replication and subsequently results in more TL attrition.

Although the associations between prenatal exposure to phthalates and newborn TL did not differ materially by trimester for many of the metabolites in our study, first-trimester exposure to ΣDEHP was related to shorter newborn TL. Early

pregnancy is generally a critical period for environmental exposures, and the developing fetus is highly vulnerable to adverse factors related to adverse birth outcomes (Mook-Kanamori et al. 2010). Unexpectedly, we also found that exposure to some phthalate metabolites during the third trimester were associated with longer newborn TL, although the association was not statistically significant. We considered that the different patterns of the associations between phthalates and newborn TL across the three trimesters might be due to the different TL dynamics during the developmental stages of the embryo and fetus. Several human and animal studies have demonstrated that cells might use different mechanisms to regulate telomere maintenance at different stages of development (Liu et al. 2007; Schatzlein et al. 2004). Furthermore, the positive associations might indicate the presence of a potential compensatory mechanism in response to phthalate exposure insults. Telomerase can synthesize telomere DNA sequences to compensate or reverse telomere erosion due to oxidative stress insults, thereby maintaining telomere length (Greider 1996; Hug and Lingner 2006; Shore and Bianchi 2009). Additional detailed studies are essential to confirm the critical time window and the potential mechanisms.

In the stratified analysis, we observed that the inverse associations between maternal urinary MECPP, MEOHP, MEHP, and ΣDEHP during the first trimester and cord blood TL were similar between male and female infants. Meanwhile, maternal exposure to MEP during the first trimester was related to shorter newborn TL only in females. The possible explanations for the sex-specific results might be the action of insulin-like growth factor 1 (IGF-1). Exposure to phthalate was reported to be inversely associated with IGF-1 (Boas et al. 2010; Wu et al. 2017). Epidemiological studies reported that IGF-1 concentration was positively associated with TL (Kaplan et al. 2009; Movérare-Skrtic et al. 2009). In addition, previous studies reported that female newborns had higher concentrations of IGF-1 than males (Geary et al. 2003; Vatten et al. 2002), which might potentially result in sex-specific effects. Moreover, evidence has suggested that increased cortisol exposure is associated with shorter TL (Choi et al. 2008; Tomiyama et al. 2012). Our previous study indicated that prenatal exposure to phthalates was positively related to the level of glucocorticoids (measured by cortisol and cortisone) in female infants but was inversely related to glucocorticoid levels in male infants (Sun et al. 2018). This might partly explain the effects of phthalates on female TL.

Our study has some strengths. The effects of prenatal phthalate exposure on newborn TL were estimated in the largest prospective study to date. In addition, the prospective cohort design and availability of urine samples from each trimester allowed us to estimate associations between newborn TL and phthalate metabolites in maternal urine samples collected at multiple time points. Moreover, we used multiple informant models to estimate the associations between maternal phthalate exposure and newborn TL according to trimester. The statistical method may be more advantageous than the separate regression models for each exposure window (Sánchez et al. 2011).

Our study also has some limitations. First, although we adjusted for some potential confounders, we cannot fully eliminate the possible residual confounding by some unobserved confounding factors. For example, the differences in individual metabolism and excretion pattern may affect urinary phthalate metabolite concentration. Second, given that phthalates are rapidly metabolized and have a short half-life, a single spot urine sample available at each trimester may not fully reflect exposure profiles at each specific window over the duration of pregnancy. Although we repeatedly measured urinary phthalate concentrations at the three trimesters, exposure measurements were still

limited due to the low ICCs, which indicated the temporal variability of phthalate concentrations over pregnancy. Third, given the high intercorrelations among several phthalate metabolites, we did not perform multi-phthalate models controlling for other individual phthalates that might be mutually confounded. To reduce the confounding by correlated phthalates, we summed all phthalate metabolites with the same chemical into one variable (i.e., DEHP). Fourth, most of the participants in the present study were Han Chinese women, and generalizability to other ethnic groups is uncertain. Furthermore, the educational level for the participants (79.0% were college or above) in the present study, which may reflect better health behaviors or higher socioeconomic status and lead to low rates of adverse pregnancy outcomes (i.e., preterm birth), may limit the generalizability of our results to other populations.

Conclusions

Our findings offer the first preliminary evidence that maternal exposure to phthalates is related to shorter TL in newborns. Such an effect may have lifetime health implications, considering that newborn TL may be critical to an individual's telomere biology and a major determinant of TL in later life (Benetos et al. 2013; Hjelmberg et al. 2015; Sabharwal et al. 2018). Future prospective investigations are warranted to further explore the effects of phthalate-related telomere shortening at birth on offspring health outcomes later in life.

Acknowledgments

We gratefully thank all the study participants in this study and the staff of the Wuhan Children's Hospital (Wuhan Maternal and Child Healthcare Hospital). This study was supported by the National Key R&D Program of China (2017YFC0212003), the National Natural Science Foundation of China (91643207, 81273083), and the Hubei Province Health and Family Planning Scientific Research Project (WJ2017Z001).

References

Arbuckle TE, Davis K, Marro L, Fisher M, Legrand M, LeBlanc A, et al. 2014. Phthalate and bisphenol A exposure among pregnant women in Canada—results from the MIREC study. *Environ Int* 68:55–65, PMID: 24709781, <https://doi.org/10.1016/j.envint.2014.02.010>.

Barker DJ. 1990. The fetal and infant origins of adult disease. *BMJ* 301(6761):1111, PMID: 2252919, <https://doi.org/10.1136/bmj.301.6761.1111>.

Bassig BA, Zhang L, Cawthon RM, Smith MT, Yin S, Li G, et al. 2014. Alterations in leukocyte telomere length in workers occupationally exposed to benzene. *Environ Mol Mutagen* 55(8):673–678, PMID: 24945723, <https://doi.org/10.1002/em.21880>.

Benetos A, Kark JD, Susser E, Kimura M, Sinnreich R, Chen W, et al. 2013. Tracking and fixed ranking of leukocyte telomere length across the adult life course. *Aging Cell* 12(4):615–621, PMID: 23601089, <https://doi.org/10.1111/accel.12086>.

Benetos A, Okuda K, Lajemi M, Kimura M, Thomas F, Skurnick J, et al. 2001. Telomere length as an indicator of biological aging: the gender effect and relation with pulse pressure and pulse wave velocity. *Hypertension* 37(2 Pt 2):381–385, PMID: 11230304, <https://doi.org/10.1161/01.HYP.37.2.381>.

Benjamin S, Masai E, Kamimura N, Takahashi K, Anderson RC, Faisal PA. 2017. Phthalates impact human health: epidemiological evidences and plausible mechanism of action. *J Hazard Mater* 340:360–383, PMID: 28800814, <https://doi.org/10.1016/j.jhazmat.2017.06.036>.

Bijnens E, Zeegers MP, Gielen M, Kicinski M, Hageman GJ, Pachon D, et al. 2015. Lower placental telomere length may be attributed to maternal residential traffic exposure; a twin study. *Environ Int* 79:1–7, PMID: 25756235, <https://doi.org/10.1016/j.envint.2015.02.008>.

Blackburn EH. 1991. Structure and function of telomeres. *Nature* 350(6319):569–573, PMID: 17081110, <https://doi.org/10.1038/350569a0>.

Boas M, Frederiksen H, Feldt-Rasmussen U, Skakkebaek NE, Hegedüs L, Hilsted L, et al. 2010. Childhood exposure to phthalates: associations with thyroid function, insulin-like growth factor I, and growth. *Environ Health Perspect* 118(10):1458–1464, PMID: 20621847, <https://doi.org/10.1289/ehp.0901331>.

Callahan CL, Pavuk M, Birnbaum LS, Ren X, Olson JR, Bonner MR. 2017. Serum polychlorinated biphenyls and leukocyte telomere length in a highly-exposed population: the Anniston Community Health Survey. *Environ Int* 108:212–220, PMID: 28886414, <https://doi.org/10.1016/j.envint.2017.08.018>.

Cantonwine DE, Meeker JD, Ferguson KK, Mukherjee B, Hauser R, McElrath TF. 2016. Urinary concentrations of bisphenol A and phthalate metabolites measured during pregnancy and risk of preeclampsia. *Environ Health Perspect* 124(10):1651–1655, PMID: 27177253, <https://doi.org/10.1289/EHP188>.

Casas M, Valvi D, Ballesteros-Gomez A, Gascon M, Fernández MF, Garcia-Esteban R, et al. 2016. Exposure to bisphenol A and phthalates during pregnancy and ultrasound measures of fetal growth in the INMA-Sabadell cohort. *Environ Health Perspect* 124(4):521–528, PMID: 26196298, <https://doi.org/10.1289/ehp.1409190>.

Cawthon RM. 2009. Telomere length measurement by a novel monochrome multiplex quantitative PCR method. *Nucleic Acids Res* 37(3):e21, PMID: 19129229, <https://doi.org/10.1093/nar/gkn1027>.

Choi J, Faure SR, Effros RB. 2008. Reduced telomerase activity in human T lymphocytes exposed to cortisol. *Brain Behav Immun* 22(4):600–605, PMID: 18222063, <https://doi.org/10.1016/j.bbi.2007.12.004>.

Diamanti-Kandarakis E, Bourguignon JP, Giudice LC, Hauser R, Prins GS, Soto AM, et al. 2009. Endocrine-disrupting chemicals: an Endocrine Society scientific statement. *Endocr Rev* 30(4):293–342, PMID: 19502515, <https://doi.org/10.1210/er.2009-0002>.

Fairlie J, Holland R, Pilkington JG, Pemberton JM, Harrington L, Nussey DH. 2016. Lifelong leukocyte telomere dynamics and survival in a free-living mammal. *Aging Cell* 15(1):140–148, PMID: 26521726, <https://doi.org/10.1111/accel.12417>.

Ferguson KK, Cantonwine DE, Rivera-González LO, Loch-Carusio R, Mukherjee B, Anzalota Del Toro LV, et al. 2014a. Urinary phthalate metabolite associations with biomarkers of inflammation and oxidative stress across pregnancy in Puerto Rico. *Environ Sci Technol* 48(12):7018–7025, PMID: 24845688, <https://doi.org/10.1021/es502076j>.

Ferguson KK, Chen YH, VanderWeele TJ, McElrath TF, Meeker JD, Mukherjee B. 2017. Mediation of the relationship between maternal phthalate exposure and preterm birth by oxidative stress with repeated measurements across pregnancy. *Environ Health Perspect* 125(3):488–494, PMID: 27352406, <https://doi.org/10.1289/EHP282>.

Ferguson KK, Loch-Carusio R, Meeker JD. 2012. Exploration of oxidative stress and inflammatory markers in relation to urinary phthalate metabolites: NHANES 1999–2006. *Environ Sci Technol* 46(1):477–485, PMID: 22085025, <https://doi.org/10.1021/es202340b>.

Ferguson KK, McElrath TF, Chen YH, Mukherjee B, Meeker JD. 2015. Urinary phthalate metabolites and biomarkers of oxidative stress in pregnant women: a repeated measures analysis. *Environ Health Perspect* 123(3):210–216, PMID: 25402001, <https://doi.org/10.1289/ehp.1307996>.

Ferguson KK, McElrath TF, Ko YA, Mukherjee B, Meeker JD. 2014b. Variability in urinary phthalate metabolite levels across pregnancy and sensitive windows of exposure for the risk of preterm birth. *Environ Int* 70:118–124, PMID: 24934852, <https://doi.org/10.1016/j.envint.2014.05.016>.

Ferguson KK, McElrath TF, Meeker JD. 2014c. Environmental phthalate exposure and preterm birth. *JAMA Pediatr* 168(1):61–67, PMID: 24247736, <https://doi.org/10.1001/jamapediatrics.2013.3699>.

Frenc RW Jr, Blackburn EH, Shannon KM. 1998. The rate of telomere sequence loss in human leukocytes varies with age. *Proc Natl Acad Sci U S A* 95(10):5607–5610, PMID: 9576930, <https://doi.org/10.1073/pnas.95.10.5607>.

Gao H, Zhu YD, Xu YY, Zhang YW, Yao HY, Sheng J, et al. 2017. Season-dependent concentrations of urinary phthalate metabolites among Chinese pregnant women: repeated measures analysis. *Environ Int* 104:110–117, PMID: 28389128, <https://doi.org/10.1016/j.envint.2017.03.021>.

Geary MP, Pringle PJ, Rodeck CH, Kingdom JC, Hindmarsh PC. 2003. Sexual dimorphism in the growth hormone and insulin-like growth factor axis at birth. *J Clin Endocrinol Metab* 88(8):3708–3714, PMID: 12915659, <https://doi.org/10.1210/jc.2002-022006>.

Greider CW. 1996. Telomere length regulation. *Annu Rev Biochem* 65:337–365, PMID: 8811183, <https://doi.org/10.1146/annurev.bi.65.070196.002005>.

Halden RU. 2010. Plastics and health risks. *Annu Rev Public Health* 31:179–194, PMID: 20070188, <https://doi.org/10.1146/annurev.publhealth.012809.103714>.

Haycock PC, Heydon EE, Kaptoge S, Butterworth AS, Thompson A, Willeit P. 2014. Leucocyte telomere length and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ* 349:g4227, PMID: 25006006, <https://doi.org/10.1136/bmj.g4227>.

Heidinger BJ, Blount JD, Boner W, Griffiths K, Metcalfe NB, Monaghan P. 2012. Telomere length in early life predicts lifespan. *Proc Natl Acad Sci U S A* 109(5):1743–1748, PMID: 22232671, <https://doi.org/10.1073/pnas.1113306109>.

Hjelmberg JB, Dalgård C, Möller S, Steenstrup T, Kimura M, Christensen K, et al. 2015. The heritability of leukocyte telomere length dynamics. *J Med Genet* 52(5):297–302, PMID: 25770094, <https://doi.org/10.1136/jmedgenet-2014-102736>.

- Holland N, Huen K, Tran V, Street K, Nguyen B, Bradman A, et al. 2016. Urinary phthalate metabolites and biomarkers of oxidative stress in a Mexican-American cohort: variability in early and late pregnancy. *Toxics* 4(1):7, PMID: 28008399, <https://doi.org/10.3390/toxics4010007>.
- Hou L, Wang S, Dou C, Zhang X, Yu Y, Zheng Y, et al. 2012. Air pollution exposure and telomere length in highly exposed subjects in Beijing, China: a repeated-measure study. *Environ Int* 48:71–77, PMID: 22871507, <https://doi.org/10.1016/j.envint.2012.06.020>.
- Houben JM, Moonen HJ, van Schooten FJ, Hageman GJ. 2008. Telomere length assessment: biomarker of chronic oxidative stress? *Free Radic Biol Med* 44(3):235–246, PMID: 18021748, <https://doi.org/10.1016/j.freeradbiomed.2007.10.001>.
- Hug N, Lingner J. 2006. Telomere length homeostasis. *Chromosoma* 115(6):413–425, PMID: 16741708, <https://doi.org/10.1007/s00412-006-0067-3>.
- Jaiswal M, LaRusso NF, Burgart LJ, Gores GJ. 2000. Inflammatory cytokines induce DNA damage and inhibit DNA repair in cholangiocarcinoma cells by a nitric oxide-dependent mechanism. *Cancer Res* 60(1):184–190, PMID: 10646872.
- Johns LE, Cooper GS, Galizia A, Meeker JD. 2015. Exposure assessment issues in epidemiology studies of phthalates. *Environ Int* 85:27–39, PMID: 26313703, <https://doi.org/10.1016/j.envint.2015.08.005>.
- Jurk D, Wilson C, Passos JF, Oakley F, Correia-Melo C, Greaves L, et al. 2014. Chronic inflammation induces telomere dysfunction and accelerates ageing in mice. *Nat Commun* 2:4172, PMID: 24960204, <https://doi.org/10.1038/ncomms5172>.
- Kaplan RC, Fitzpatrick AL, Pollak MN, Gardner JP, Jenny NS, McGinn AP, et al. 2009. Insulin-like growth factors and leukocyte telomere length: the Cardiovascular Health Study. *J Gerontol A Biol Sci Med Sci* 64(11):1103–1106, PMID: 19349587, <https://doi.org/10.1093/gerona/glp036>.
- Kaufman JS, MacLehose RF. 2013. Which of these things is not like the others? *Cancer* 119(24):4216–4222, PMID: 24022386, <https://doi.org/10.1002/cncr.28359>.
- Kawanishi S, Oikawa S. 2004. Mechanism of telomere shortening by oxidative stress. *Ann N Y Acad Sci* 1019(1):278–284, PMID: 15247029, <https://doi.org/10.1196/annals.1297.047>.
- Koch HM, Calafat AM. 2009. Human body burdens of chemicals used in plastic manufacture. *Philos Trans R Soc Lond B Biol Sci* 364(1526):2063–2078, PMID: 19528056, <https://doi.org/10.1098/rstb.2008.0208>.
- Latini G, De Felice C, Presta G, Del Vecchio A, Paris I, Ruggieri F, et al. 2003. *In utero* exposure to di-(2-ethylhexyl)phthalate and duration of human pregnancy. *Environ Health Perspect* 111(14):1783–1785, PMID: 14594632, <https://doi.org/10.1289/ehp.6202>.
- Li H, Engström K, Vahter M, Broberg K. 2012. Arsenic exposure through drinking water is associated with longer telomeres in peripheral blood. *Chem Res Toxicol* 25(11):2333–2339, PMID: 22917110, <https://doi.org/10.1021/tx300222t>.
- Liu H, Chen Q, Lei L, Zhou W, Huang L, Zhang J, et al. 2018. Prenatal exposure to perfluoroalkyl and polyfluoroalkyl substances affects leukocyte telomere length in female newborns. *Environ Pollut* 235:446–452, PMID: 29310088, <https://doi.org/10.1016/j.envpol.2017.12.095>.
- Liu L, Bailey SM, Okuka M, Muñoz P, Li C, Zhou L, et al. 2007. Telomere lengthening early in development. *Nat Cell Biol* 9(12):1436–1441, PMID: 17982445, <https://doi.org/10.1038/ncb1664>.
- Marchetto NM, Glynn RA, Ferry ML, Ostojic M, Wolff SM, Yao R, et al. 2016. Prenatal stress and newborn telomere length. *Am J Obstet Gynecol* 215(94):e1–e8, PMID: 26829506, <https://doi.org/10.1016/j.ajog.2016.01.177>.
- Martens DS, Cox B, Janssen BG, Clemente DBP, Gasparrini A, Vanpoucke C, et al. 2017. Prenatal air pollution and newborns' predisposition to accelerated biological aging. *JAMA Pediatr* 171(12):1160–1167, PMID: 29049509, <https://doi.org/10.1001/jamapediatrics.2017.3024>.
- Martens DS, Plusquin M, Gyselaers W, De Vivo I, Nawrot TS. 2016. Maternal pre-pregnancy body mass index and newborn telomere length. *BMC Med* 14(1):148, PMID: 27751173, <https://doi.org/10.1186/s12916-016-0689-0>.
- McCracken J, Baccarelli A, Hoxha M, Dioni L, Melly S, Coull B, et al. 2010. Annual ambient black carbon associated with shorter telomeres in elderly men: Veterans Affairs Normative Aging Study. *Environ Health Perspect* 118(11):1564–1570, PMID: 21465749, <https://doi.org/10.1289/ehp.0901831>.
- Mirzakhani H, De Vivo I, Leeder JS, Gaedigk R, Vyhldal CA, Weiss ST, et al. 2017. Early pregnancy intrauterine fetal exposure to maternal smoking and impact on fetal telomere length. *Eur J Obstet Gynecol Reprod Biol* 218:27–32, PMID: 28926727, <https://doi.org/10.1016/j.ejogrb.2017.09.013>.
- Mitro SD, Birnbaum LS, Needham BL, Zota AR. 2016. Cross-sectional associations between exposure to persistent organic pollutants and leukocyte telomere length among U.S. adults in NHANES, 2001–2002. *Environ Health Perspect* 124(5):651–658, PMID: 26452299, <https://doi.org/10.1289/ehp.1510187>.
- Mons U, Müezziner A, Schöttker B, Dieffenbach AK, Butterbach K, Schick M, et al. 2017. Leukocyte telomere length and all-cause, cardiovascular disease, and cancer mortality: results from individual-participant-data meta-analysis of 2 large prospective cohort studies. *Am J Epidemiol* 185(12):1317–1326, PMID: 28459963, <https://doi.org/10.1093/aje/kww210>.
- Mook-Kanamori DO, Steegers EA, Eilers PH, Raat H, Hofman A, Jaddoe VW. 2010. Risk factors and outcomes associated with first-trimester fetal growth restriction. *JAMA* 303(6):527–534, PMID: 20145229, <https://doi.org/10.1001/jama.2010.78>.
- Moreno-Palomo J, Creus A, Marcos R, Hernández A. 2014. Genomic instability in newborn with short telomeres. *PLoS One* 9(3):e91753, PMID: 24622247, <https://doi.org/10.1371/journal.pone.0091753>.
- Mose T, Mortensen GK, Hedegaard M, Knudsen LE. 2007. Phthalate monoesters in perfusate from a dual placenta perfusion system, the placenta tissue and umbilical cord blood. *Reprod Toxicol* 23(1):83–91, PMID: 17049806, <https://doi.org/10.1016/j.reprotox.2006.08.006>.
- Movérare-Skrtc S, Svensson J, Karlsson MK, Orwoll E, Ljunggren O, Mellström D, et al. 2009. Serum insulin-like growth factor-I concentration is associated with leukocyte telomere length in a population-based cohort of elderly men. *J Clin Endocrinol Metab* 94(12):5078–5084, PMID: 19846733, <https://doi.org/10.1210/jc.2009-1450>.
- Mu D, Gao F, Fan Z, Shen H, Peng H, Hu J. 2015. Levels of phthalate metabolites in urine of pregnant women and risk of clinical pregnancy loss. *Environ Sci Technol* 49(17):10651–10657, PMID: 26251123, <https://doi.org/10.1021/acs.est.5b02617>.
- Myrstadakis A, Fthenou E, Balaska E, Vakinti M, Kogevinas M, Stephanou EG. 2015. Phthalate esters, parabens and bisphenol-A exposure among mothers and their children in Greece (Rhea cohort). *Environ Int* 83:1–10, PMID: 26072145, <https://doi.org/10.1016/j.envint.2015.05.014>.
- Perera F, Lin CJ, Qu L, Tang D. 2018. Shorter telomere length in cord blood associated with prenatal air pollution exposure: benefits of intervention. *Environ Int* 113:335–340, PMID: 29395277, <https://doi.org/10.1016/j.envint.2018.01.005>.
- Philippat C, Mortamais M, Chevrier C, Petit C, Calafat AM, Ye X, et al. 2012. Exposure to phthalates and phenols during pregnancy and offspring size at birth. *Environ Health Perspect* 120(3):464–470, PMID: 21900077, <https://doi.org/10.1289/ehp.1103634>.
- Pieters N, Janssen BG, Dewitte H, Cox B, Cuypers A, Lefebvre W, et al. 2016. Biomolecular markers within the core axis of aging and particulate air pollution exposure in the elderly: a cross-sectional study. *Environ Health Perspect* 124(7):943–950, PMID: 26672058, <https://doi.org/10.1289/ehp.1509728>.
- Qu F, Li R, He X, Li Q, Xie S, Gong L, et al. 2015. Short telomere length in peripheral blood leukocyte predicts poor prognosis and indicates an immunosuppressive phenotype in gastric cancer patients. *Mol Oncol* 9(3):727–739, PMID: 25515040, <https://doi.org/10.1016/j.molonc.2014.11.008>.
- Rudel RA, Gray JM, Engel CL, Rawsthorne TW, Dodson RE, Ackerman JM, et al. 2011. Food packaging and bisphenol A and bis(2-ethylhexyl) phthalate exposure: findings from a dietary intervention. *Environ Health Perspect* 119(7):914–920, PMID: 21450549, <https://doi.org/10.1289/ehp.1003170>.
- Sabharwal S, Verhulst S, Guirguis G, Kark JD, Labat C, Roche NE, et al. 2018. Telomere length dynamics in early life: the blood-and-muscle model. *FASEB J* 32(1):529–534, PMID: 28855279, <https://doi.org/10.1096/fj.201700630R>.
- Sánchez BN, Hu H, Litman HJ, Téllez-Rojo MM. 2011. Statistical methods to study timing of vulnerability with sparsely sampled data on environmental toxicants. *Environ Health Perspect* 119(3):409–415, PMID: 21362588, <https://doi.org/10.1289/ehp.1002453>.
- Schaetzlein S, Lucas-Hahn A, Lemme E, Kues WA, Dorsch M, Manns MP, et al. 2004. Telomere length is reset during early mammalian embryogenesis. *Proc Natl Acad Sci U S A* 101(21):8034–8038, PMID: 15148368, <https://doi.org/10.1073/pnas.0402400101>.
- Schettler T. 2006. Human exposure to phthalates via consumer products. *Int J Androl* 29(1):134–139, PMID: 16466533, <https://doi.org/10.1111/j.1365-2605.2005.00567.x>.
- Scinicariello F, Feroe AG, Attanasio R. 2016. Urinary phthalates and leukocyte telomere length: an analysis of NHANES 1999–2002. *EBioMedicine* 6:96–102, PMID: 27211552, <https://doi.org/10.1016/j.ebiom.2016.02.027>.
- Shore D, Bianchi A. 2009. Telomere length regulation: coupling DNA end processing to feedback regulation of telomerase. *EMBO J* 28(16):2309–2322, PMID: 19629031, <https://doi.org/10.1038/emboj.2009.195>.
- Soomro MH, Baiz N, Philippat C, Vernet C, Siroux V, Nichole Maesano C, et al. 2018. Prenatal exposure to phthalates and the development of eczema phenotypes in male children: results from the EDEN mother–child cohort study. *Environ Health Perspect* 126(2):027002, PMID: 29398652, <https://doi.org/10.1289/EHP1829>.
- Sun X, Li J, Jin S, Li Y, Liu W, Zhao H, et al. 2018. Associations between repeated measures of maternal urinary phthalate metabolites during pregnancy and cord blood glucocorticoids. *Environ Int* 121(Pt 1):471–479, PMID: 30278310, <https://doi.org/10.1016/j.envint.2018.09.037>.
- Tefre de Renzy-Martin K, Frederiksen H, Christensen JS, Boye Kyhl H, Andersson AM, Husby S, et al. 2014. Current exposure of 200 pregnant Danish women to phthalates, parabens and phenols. *Reproduction* 147(4):443–453, PMID: 24282315, <https://doi.org/10.1530/REP-13-0461>.

- Téllez-Rojo MM, Cantoral A, Cantonwine DE, Schnaas L, Peterson K, Hu H, et al. 2013. Prenatal urinary phthalate metabolites levels and neurodevelopment in children at two and three years of age. *Sci Total Environ* 461-462:386–390, PMID: [23747553](#), <https://doi.org/10.1016/j.scitotenv.2013.05.021>.
- Tomiya AJ, O'Donovan A, Lin J, Puterman E, Lazaro A, Chan J, et al. 2012. Does cellular aging relate to patterns of allostasis? An examination of basal and stress reactive HPA axis activity and telomere length. *Physiol Behav* 106(1):40–45, PMID: [22138440](#), <https://doi.org/10.1016/j.physbeh.2011.11.016>.
- Turner KJ, Vasu V, Griffin DK. 2019. Telomere biology and human phenotype. *Cells* 8(1):E73, PMID: [30669451](#), <https://doi.org/10.3390/cells8010073>.
- Vardavas CI, Hohmann C, Patelarou E, Martínez D, Henderson AJ, Granell R, et al. 2016. The independent role of prenatal and postnatal exposure to active and passive smoking on the development of early wheeze in children. *Eur Respir J* 48(1):115–124, PMID: [26965294](#), <https://doi.org/10.1183/13993003.01016-2015>.
- Vatten LJ, Nilsen ST, Ødegård RA, Romundstad PR, Austgulen R. 2002. Insulin-like growth factor I and leptin in umbilical cord plasma and infant birth size at term. *Pediatrics* 109(6):1131–1135, PMID: [12042554](#), <https://doi.org/10.1542/peds.109.6.1131>.
- Wai KM, Umezaki M, Kosaka S, Mar O, Umemura M, Fillman T, et al. 2018. Impact of prenatal heavy metal exposure on newborn leucocyte telomere length: a birth-cohort study. *Environ Pollut* 243(Pt B):1414–1421, PMID: [30278415](#), <https://doi.org/10.1016/j.envpol.2018.09.090>.
- Watkins DJ, Sánchez BN, Téllez-Rojo MM, Lee JM, Mercado-García A, Blank-Goldenberg C, et al. 2017a. Impact of phthalate and BPA exposure during in utero windows of susceptibility on reproductive hormones and sexual maturation in peripubertal males. *Environ Health* 16(1):69, PMID: [28637469](#), <https://doi.org/10.1186/s12940-017-0278-5>.
- Watkins DJ, Sánchez BN, Téllez-Rojo MM, Lee JM, Mercado-García A, Blank-Goldenberg C, et al. 2017b. Phthalate and bisphenol A exposure during in utero windows of susceptibility in relation to reproductive hormones and pubertal development in girls. *Environ Res* 159:143–151, PMID: [28800472](#), <https://doi.org/10.1016/j.envres.2017.07.051>.
- Willeit P, Willeit J, Mayr A, Weger S, Oberhollenzer F, Brandstätter A, et al. 2010. Telomere length and risk of incident cancer and cancer mortality. *JAMA* 304(1):69–75, PMID: [20606151](#), <https://doi.org/10.1001/jama.2010.897>.
- Wittassek M, Koch HM, Angerer J, Brüning T. 2011. Assessing exposure to phthalates – the human biomonitoring approach. *Mol Nutr Food Res* 55(1):7–31, PMID: [20564479](#), <https://doi.org/10.1002/mnfr.201000121>.
- Wormuth M, Scheringer M, Vollenweider M, Hungerbühler K. 2006. What are the sources of exposure to eight frequently used phthalic acid esters in Europeans? *Risk Anal* 26(3):803–824, PMID: [16834635](#), <https://doi.org/10.1111/j.1539-6924.2006.00770.x>.
- Wu W, Zhou F, Wang Y, Ning Y, Yang JY, Zhou YK. 2017. Exposure to phthalates in children aged 5–7 years: associations with thyroid function and insulin-like growth factors. *Sci Total Environ* 579:950–956, PMID: [27884522](#), <https://doi.org/10.1016/j.scitotenv.2016.06.146>.
- Wu Y, Liu Y, Ni N, Bao B, Zhang C, Lu L. 2012. High lead exposure is associated with telomere length shortening in Chinese battery manufacturing plant workers. *Occup Environ Med* 69(8):557–563, PMID: [22539657](#), <https://doi.org/10.1136/oemed-2011-100478>.
- Zhang YW, Gao H, Mao LJ, Tao XY, Ge X, Huang K, et al. 2018. Effects of the phthalate exposure during three gestation periods on birth weight and their gender differences: a birth cohort study in China. *Sci Total Environ* 613–614:1573–1578, PMID: [28886917](#), <https://doi.org/10.1016/j.scitotenv.2017.08.319>.
- Zhao J, Miao K, Wang H, Ding H, Wang DW. 2013. Association between telomere length and type 2 diabetes mellitus: a meta-analysis. *PLoS One* 8(11):e79993, PMID: [24278229](#), <https://doi.org/10.1371/journal.pone.0079993>.
- Zhu Y, Wan Y, Li Y, Zhang B, Zhou A, Cai Z, et al. 2016. Free and total urinary phthalate metabolite concentrations among pregnant women from the Healthy Baby Cohort (HBC), China. *Environ Int* 88:67–73, PMID: [26722670](#), <https://doi.org/10.1016/j.envint.2015.12.004>.
- Zhu Y, Wan Y, Zhang B, Zhou A, Huo W, Wu C, et al. 2018. Relationship between maternal phthalate exposure and offspring size at birth. *Sci Total Environ* 612:1072–1078, PMID: [28892847](#), <https://doi.org/10.1016/j.scitotenv.2017.08.207>.
- Zota AR, Needham BL, Blackburn EH, Lin J, Park SK, Rehkopf DH, et al. 2015. Associations of cadmium and lead exposure with leukocyte telomere length: findings from National Health and Nutrition Examination Survey, 1999–2002. *Am J Epidemiol* 181(2):127–136, PMID: [25504027](#), <https://doi.org/10.1093/aje/kwu293>.